

Symmetrized Schrödinger Equation and General Complex Solution *

YiHuan Wei^{1,2}

¹ Institute of Theoretical Physics, Chinese Academy of Science, Beijing 100080, China

² Department of Physics, Bohai University, Jinzhou 121000, Liaoning, China

February 1, 2008

Abstract

We suggest the symmetrized Schrödinger equation and propose a general complex solution which is characterized by the imaginary units i and ϵ . This symmetrized Schrödinger equation appears some interesting features.

I Introduction

Quantum physics was rapidly developed in the first three decades of last century (for a time list of some important findings, see Ref.[1]). The fundamental framework of quantum mechanics was set up in 1920's [2, 3], (for a review, see Ref. [4]). The interference of two noninteracting beams of "particles" and the diffraction of single beam of "particles" enforced us to admit its wave property. The wavefunction describing the states of the physical system evolves according to the time-dependent Schrödinger equation. This wave equation, which contains the first order derivatives of the wavefunction in time, can be found by the classical one by the replacements of some classical quantities by the corresponding operators. However, the noncommutativity of general coordinate operator $\hat{q}$ and momentum operator $\hat{p}$

*Liaoning Province Educational Committee (9808111084).

leads to ambiguities in the definition of the operator $\hat{H}$, the wave equation (the Schrödinger equation) is thus taken as a fundamental equation independent on a underlying classical picture [5]. Within quantum mechanics, the wavefunction is not a observable and what it describes is not the wave propagating in usual three-dimensional space but a representative element of the state of the system. The state is only a ray (or vector) in abstract *Hilbert* space, without any classical counterpart.

In quantum mechanics, one of the postulates states that to any self-consistent and well-defined observable in physics corresponds to an operator, the value of the observable corresponds to the eigenvalue (expectation value) of the operator. We will use the term expectation value, instead of eigenvalue, one can see the former will fit more to our discussions since we will deal with a symmetrized Schrödinger equation. The operator corresponding to the energy is the Hamiltonian $\hat{H}$. For a single particle of the mass μ , in a potential $V(\mathbf{x})$, $\hat{H} = \frac{\hat{\mathbf{p}}^2}{2\mu} + V(\mathbf{x})$ with $\hat{\mathbf{p}} = -i\hbar\nabla$. The expectation value equation, i.e, the time-dependent Schrödinger equation yields the possible energy which the particle may have. For a free particle the expectation value of the energy is $E = \frac{\mathbf{p}^2}{2\mu}$ with $\mathbf{p}$ the expectation value of the momentum. This is completely consistent with the classical relationship between kinetic and momentum.

As some preparations, let us introduce some knowledge of the complex numbers. The hyperbolic complex numbers may be generated by the hyperbolic imaginary unit ϵ satisfying $\epsilon^2 = 1$. Similarly to the ordinary complex numbers, a hyperbolic complex number takes the form [6]

$$Z_h = x + \epsilon y, \tag{1}$$

where x and y are two real numbers, called the real and imaginary parts of the hyperbolic complex number Z_h . The exponent form of the hyperbolic complex number Z_h is given as

follows

$$Z_h = e^{\epsilon\theta} = \cosh \theta + \epsilon \sinh \theta, \quad (2)$$

where θ is a real parameter. For a hyperbolic complex number defined by (1), the conjugate is

$$Z_h^* = x - \epsilon y. \quad (3)$$

The square norm of a hyperbolic complex number is defined as $|Z_h|^2 = Z_h^* Z_h = x^2 - y^2$. The set of hyperbolic complex numbers has the same algebraic laws as the one of complex numbers. It has been applied in many fields, such as, gravitational fields [8], quantum group [7], quantum mechanics [9], *etc.* Recently, we propose so-called η -complex number which has the two specific η -values ∓ 1 corresponding to the usual complex and hyperbolic complex numbers, respectively [10].

In this paper, we suggest the symmetrized Schrödinger equation. This equation doesn't only contain the usual Schrödinger equation but also shows some interesting features. It allows such solutions that are characterized by the two imaginary units i and ϵ . The three equations derived from it have an excellent characteristic that automatically gives the constraint on the function $\mathcal{A} = \mathcal{A}(x)$ and admits a more general function $\mathcal{S} = \mathcal{S}(x, t)$. This paper is organized as follows. In Sec.II we discuss in detailed the equations derived from the symmetrized Schrödinger equation and gives some special solutions. In Sec.III we give a simple comparison between the usual Schrödinger equation and the symmetrized Schrödinger equation, and analyze the relation between kinetic energy and momentum of free particle in our framework.

II The Symmetrized Schrödinger Equation

The time-dependent Schrödinger equation plays a fundamental role in quantum mechanics.

It takes the following form

$$i\frac{\partial}{\partial t}\psi = \hat{H}\psi, \quad (4)$$

where for simplicity we have used the unit in which $\hbar = 1$, in which the imaginary unit i takes an important place. Eq.(4) has the complex solution $\psi = \mathcal{A}e^{i\mathcal{S}}$ or the hyperbolic complex solution $\psi = \mathcal{A}e^{\epsilon\mathcal{S}}$ (see Ref.[9]), where $\mathcal{A} = \mathcal{A}(x, t)$ and $\mathcal{S} = \mathcal{S}(x, t)$ are the amplitude and phase, respectively.

Naturally, the Schrödinger equation may be reformulated as

$$\frac{1}{2}(i\frac{\partial}{\partial t}\psi + \frac{\partial}{\partial t}\psi i) = \hat{H}\psi. \quad (5)$$

Here, let us call this equation the symmetrized Schrödinger Equation. For the equation (5) we present the following complex solution of the form

$$\psi = \mathcal{A}e^{(\delta i + \sigma\epsilon)\mathcal{S}}, \quad (6)$$

which comprises i and ϵ , where δ and σ are two constants, $\mathcal{A}$ and $\mathcal{S}$ are still called the amplitude and phase, respectively. Noting that (see Ref.[7])

$$\{i, \epsilon\} = i\epsilon + \epsilon i = 0, \quad (7)$$

by expanding ψ given by Eq.(6) in $\delta i + \sigma\epsilon$, then we have

$$\psi = \mathcal{A}[c(\mathcal{S}; \delta, \sigma) + (\delta i + \sigma\epsilon)s(\mathcal{S}; \delta, \sigma)], \quad (8)$$

where

$$c(\mathcal{S}; \delta, \sigma) = \sum_{n=0}^{\infty} \frac{(\sigma^2 - \delta^2)^n \mathcal{S}^{2n}}{(2n)!}, \quad s(\mathcal{S}; \delta, \sigma) = \sum_{n=0}^{\infty} \frac{(\sigma^2 - \delta^2)^n \mathcal{S}^{2n+1}}{(2n+1)!}, \quad (9)$$

which will reduce to $c(\mathcal{S}; \delta, \sigma) = \cos \mathcal{S}$ and $s(\mathcal{S}; \delta, \sigma) = \sin \mathcal{S}$ for $\sigma^2 - \delta^2 = -1$, $c(\mathcal{S}; \delta, \sigma) = \cosh \mathcal{S}$ and $s(\mathcal{S}; \delta, \sigma) = \sinh \mathcal{S}$ for $\sigma^2 - \delta^2 = 1$. One can find the following relation between the function $c(\mathcal{S}; \delta, \sigma)$ and $s(\mathcal{S}; \delta, \sigma)$

$$c^2(\mathcal{S}; \delta, \sigma) - (\sigma^2 - \delta^2)s^2(\mathcal{S}; \delta, \sigma) = 1, \quad (10)$$

which will give $\cos^2 \mathcal{S} + \sin^2 \mathcal{S} = 1$ and $\cosh^2 \mathcal{S} - \sinh^2 \mathcal{S} = 1$ for $\sigma^2 - \delta^2 = -1, 1$, respectively. In fact, Eqs.(8)-(10) has been given in Ref.[10]. From the representations of i and ϵ [7], one can have

$$\delta i + \sigma \epsilon = \delta \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} + \sigma \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \sigma - \delta \\ \delta + \sigma & 0 \end{pmatrix}. \quad (11)$$

Letting $\eta = \sigma^2 - \delta^2$, then Eq.(11) gives a representation [7] of i_η with $i_\eta^2 = \eta$. The wavefunction (6) can take the form $\psi = \mathcal{A}e^{i_\eta \mathcal{S}}$ and therefore some calculations related to the derivative of the functions $s(\mathcal{S}; \delta, \sigma)$ and $c(\mathcal{S}; \delta, \sigma)$ or $\psi = \mathcal{A}e^{(\delta i + \sigma \epsilon)\mathcal{S}}$ are directly applicable, here.

According to Ref.[10], we have

$$\frac{\partial}{\partial q}s(\mathcal{S}; \delta, \sigma) = c(\mathcal{S}; \delta, \sigma)\frac{\partial \mathcal{S}}{\partial q}, \quad \frac{\partial}{\partial q}c(\mathcal{S}; \delta, \sigma) = (\sigma^2 - \delta^2)s(\mathcal{S}; \delta, \sigma)\frac{\partial \mathcal{S}}{\partial q}, \quad (12)$$

where $q = t, x$, or for the wavefunction $\psi = \mathcal{A}e^{(\delta i + \sigma \epsilon)\mathcal{S}}$,

$$\frac{\partial}{\partial q}\psi = \psi \frac{\partial \ln \mathcal{A}}{\partial q} + (\delta i + \sigma \epsilon)\psi \frac{\partial \mathcal{S}}{\partial q}. \quad (13)$$

As an illustrative example of Eq.(6), we will consider the case of free particle of the mass μ in one-dimensional space. In this case, $\mathcal{A} = \text{const}$, $\mathcal{S} = px - Et$ with p and E two parameters, the right side (RS) and left side (LS) of Eq.(5) are

$$RS = (\delta^2 - \sigma^2)\frac{p^2}{2\mu}\psi, \quad LS = \delta E\psi. \quad (14)$$

Equating RS to LS in (14) gives

$$E = \frac{\delta^2 - \sigma^2}{\delta} \frac{p^2}{2\mu}. \quad (15)$$

This equation shows the relation between the parameter E and p . When $\frac{\delta^2 - \sigma^2}{\delta} = 1$ (two special solutions being $(\delta = 1, \sigma = 0)$ and $(\delta = -1, \sigma = \pm\sqrt{2})$), it is compatible with the classical relation between kinetic energy and momentum.

In order to derive the more general equations stemming from the symmetrized Schrödinger equation with the wavefunction (6), let us return to the case of three-dimensional spaces. Writing $\mathcal{A} = \mathcal{A}(\mathbf{x}, t)$ and $\mathcal{S} = \mathcal{S}(\mathbf{x}, t)$, where $\mathbf{x} = (x_i, i = 1, 2, 3)$, and noting (7), then the symmetrized Schrödinger equation (5) yields

$$\delta \mathcal{A} \dot{\mathcal{S}} + i \dot{\mathcal{A}} = -\frac{1}{2\mu} [\nabla^2 \mathcal{A} + (\delta i + \sigma \epsilon)(\mathcal{A} \nabla^2 \mathcal{S} + 2 \nabla \mathcal{A} \cdot \nabla \mathcal{S}) + (\delta i + \sigma \epsilon)^2 \mathcal{A} \nabla \mathcal{S} \cdot \nabla \mathcal{S}] + V \mathcal{A}. \quad (16)$$

where $V = V(\mathbf{x})$ is a potential function. Since 1, i and ϵ are three independent algebraic generators, (16) will yield the three independent equations

$$\nabla^2 \mathcal{A} + (\sigma^2 - \delta^2) \mathcal{A} \nabla \mathcal{S} \cdot \nabla \mathcal{S} + 2\mu \delta \mathcal{A} \dot{\mathcal{S}} - 2\mu V \mathcal{A} = 0, \quad (17)$$

$$\dot{\mathcal{A}} = -\frac{1}{2\mu} \delta (\mathcal{A} \nabla^2 \mathcal{S} + 2 \nabla \mathcal{A} \cdot \nabla \mathcal{S}), \quad (18)$$

$$\mathcal{A} \nabla^2 \mathcal{S} + 2 \nabla \mathcal{A} \cdot \nabla \mathcal{S} = 0, \quad (19)$$

where $\dot{\mathcal{A}}$ denotes the derivatives with respect to time. The last of these three equations is the same as (12b) in Ref.[9], and the first one will reduce to (12a) when $\sigma^2 - \delta^2 = 1$ for $\dot{\mathcal{S}} = 0$. There is also an additional equation (18), which prescribes the derivative of $\mathcal{A}$ with respect to time. Clearly, combining the second and third one leads to $\dot{\mathcal{A}} = 0$.

In the case of one-dimensional space, the system of equations (17)-(19) is reduced to

$$\mathcal{A}'' + (\sigma^2 - \delta^2) \mathcal{A} \mathcal{S}'^2 + 2\mu \delta \mathcal{A} \dot{\mathcal{S}} - 2\mu V \mathcal{A} = 0, \quad (20)$$

$$\mathcal{A} \mathcal{S}'' + 2 \mathcal{A}' \mathcal{S}' = 0, \quad (21)$$

with $\mathcal{A} = \mathcal{A}(x)$ which is the consequence of Eqs.(18) and (19). For the cases of $\delta = 1$, $\sigma = 0$ and $\sigma = 1$, $\delta = 0$, Eq.(20) corresponds to the usual complex and hyperbolic complex

situations [9], respectively. Here, we will focus on the third special case $\sigma^2 - \delta^2 = 0$. In this case, Eq.(20) is further reduced to

$$\mathcal{A}'' + 2\mu\delta\mathcal{A}\dot{\mathcal{S}} - 2\mu V\mathcal{A} = 0. \quad (22)$$

First, let $\mathcal{S}$ be only a function of space $\mathcal{S} = \mathcal{S}(x)$. In this case, Eq.(22) is

$$\mathcal{A}'' - 2\mu V\mathcal{A} = 0. \quad (23)$$

Although this equation take a simple form and can familiar to us we still want to show some representative examples, such as, the constant potential, the inverse quadratic potential, the quadratic potential, since we are considering a special case of $\sigma^2 - \delta^2 = 0$.

For the constant potential $V = -\eta V_0$ with $V_0 > 0$ and $\eta = \pm 1$, the solution of Eqs.(23) and (21) is given by

$$\mathcal{A} = C_1 c(kx; \eta) + C_2, \quad \mathcal{S} = B_1 \int \mathcal{A}^{-2} dx + B_2, \quad (24)$$

with $k = (2\mu V_0)^{1/2}$, where C_1 , C_2 , B_1 and B_2 are integrating constants. For the $C_2 = 0$ case, $\mathcal{S} = B_1 C_1^{-2} k^{-1} \frac{s(kx; \eta)}{c(kx; \eta)} + B_2$.

For the potential $V = V_0 x^{-2}$ with V_0 a constant, the solution is

$$\mathcal{A} = Cx^n, \quad \mathcal{S} = B_1 x^{-2n+1} + B_2, \quad (25)$$

where C is a constant and n is given by $n^2 - n - 2\mu V_0 = 0$ with $V_0 > -\frac{1}{8\mu}$.

For the potential $V = \frac{1}{2\mu} \sum_{n=0}^2 V_n x^n$ with V_n some constants, $V_2 > 0$ and V_0 being determined by $V_0 = \frac{V_1^2}{4V_2} \pm \sqrt{V_2}$, the solution is obtained by

$$\mathcal{A} = e^f, \quad f = \sum_{m=0}^2 C_m x^m, \quad \mathcal{S} = B_1 \int \mathcal{A}^{-2} dx + B_2 \quad (26)$$

where $C_2 = \pm \frac{1}{2} \sqrt{V_2}$, $C_1 = \pm \frac{V_1}{2\sqrt{V_2}}$ and C_0 is an arbitrary constant. Performing the integral of $\mathcal{S}$ in (26) gives $\mathcal{S} = -\frac{\sqrt{\pi} B_1}{2\sqrt{2C_2}} e^{\frac{C_1^2}{2C_2} - 2C_0} \text{erf}(\sqrt{2C_2}(x + \frac{C_1}{2C_2})) + B_2$ with $C_2 > 0$, where $\text{erf}(x)$ is the error function which takes on values on the range -1 and 1 .

Subsequently, we consider the situation of $\mathcal{S} = \mathcal{S}(t)$. In this case, Eq.(21) is automatically satisfied, and we need only deal with Eq.(22). Letting $\dot{\mathcal{S}} = E = E(t)$, then from (22) we have

$$\mathcal{A}'' + 2\mu(\delta E - V)\mathcal{A} = 0. \quad (27)$$

Noting that $\mathcal{A} = \mathcal{A}(x)$, one should require that $\delta E - V$ is only a function of space. Whether E is time-dependent or not will depend on whether a potential V is or not. Defining $\tilde{E} = \tilde{E}(x) = -\delta E + V$, then Eq.(27) takes precisely the form of Eq.(23) and therefore has the solutions that $\mathcal{A}$'s are given by Eqs.(24)-(26) for $\tilde{E} = -\eta V_0$, $V_0 x^{-2}$, $\sum_{n=0}^2 V_n x^n$, which correspond to the potentials $V = -\eta V_0 - \delta E$, $V_0 x^{-2} - \delta E$, $\sum_{n=0}^2 V_n x^n - \delta E$, respectively.

Now, let us turn to Eqs. (20) and (21) with an arbitrary $\sigma^2 - \delta^2$. We have a concise method of solving these two equations. In fact, there always is the following relation between $\mathcal{S}$ and $\mathcal{A}$

$$\mathcal{S} = B_1 \int \mathcal{A}^{-2} dx + B_2 \quad (28)$$

or $\mathcal{S}' = B_1 \mathcal{A}^{-2}$. For the $\dot{\mathcal{S}} = 0$ case, putting Eq.(28) into (20) yields

$$\mathcal{A}'' + (\sigma^2 - \delta^2)B_1^2 \mathcal{A}^{-3} - 2\mu V \mathcal{A} = 0. \quad (29)$$

It is easy to check that Eq.(29) have such solutions that take the forms of these given in Table I in [9].

Here, we show a solution of Eq.(29) as follows

$$\mathcal{A} = Bx^{-\frac{n-1}{2}} e^{k_n x^n}, \quad \mathcal{S} = -\frac{B_1}{2nk_n B^2} e^{-2k_n x^n} + B_2, \quad (30)$$

with B and k_n some constants, which corresponds to the following potential

$$V = \frac{1}{2\mu} \left[\frac{n^2 - 1}{4} x^{-2} + n^2 k_n^2 x^{2(n-1)} + (\sigma^2 - \delta^2) B_1^2 B^{-4} x^{2(n-1)} e^{-4k_n x^n} \right]. \quad (31)$$

For $\sigma^2 - \delta^2 = 1$ and $n = 1$, the solution (30) give the third solution in Table I, where we find the constant C appearing in the solution with $V = Ce^{4kx}$ [9] should be constrained to the range $C \geq 0$.

III Discussions

The original Schrödinger equation has no such a solution as (6) with nonzero δ and σ , thus the symmetrized Schrödinger equation should be a nontrivial generalization. Our solution (6) has taken a more general form, which contains both the usual complex solution and hyperbolic complex solution [9] and the other special case which comprises two complex numbers, i.e., the solution given in terms of the mixture of two complex numbers.

Here, let us proceed with our analysis of the case $\sigma^2 - \delta^2 = 0$. According to (9), for $\sigma^2 - \delta^2 = 0$ we have $s(\mathcal{S}; \delta, \sigma) = \mathcal{S}$ and $c(\mathcal{S}; \delta, \sigma) = 1$ and therefore the wavefunction (8) takes the form $\psi = \mathcal{A}(1 + i_0\mathcal{S})$ or equivalently $\psi = \mathcal{A}e^{i_0\mathcal{S}}$, where $i_0 \neq 0$ the representation of which may be given from (11) by letting $\sigma = \pm\delta$. The conjugate of ψ is $\psi^* = \mathcal{A}e^{-i_0\mathcal{S}} = \mathcal{A}(1 - i_0\mathcal{S})$ and the square norm $|\psi|^2 = \psi^*\psi = \mathcal{A}^2$, the definitions of the conjugate and norm still holding valid in this special case.

For $\delta = 0$ and $\sigma = 1$, the solution (6) may be formally reduced to that of [9] but it allows for the function $\mathcal{S}$ to take a more general form $\mathcal{S} = \mathcal{S}(x, t)$. For the constant $\mathcal{A}$ and $\mathcal{S} = \mathcal{S}(x, t)$, Eq.(21) will yield $\mathcal{S} = \mathcal{S}_1x + \mathcal{S}_2$, where $\mathcal{S}_1 = \mathcal{S}_1(t)$, $\mathcal{S}_2 = \mathcal{S}_2(t)$. Putting it into Eq.(20) gives

$$(\sigma^2 - \delta^2)\mathcal{S}_1^2 + 2\mu\delta(\dot{\mathcal{S}}_1x + \dot{\mathcal{S}}_2) - 2\mu V = 0. \quad (32)$$

Letting $\mathcal{S}_1 = p = p_0 + kt$ with p_0 and k two constants and $\mathcal{S}_2 = Et$ with E also a constant, then (32) reduces to $E + \dot{E}t + kx - \delta^{-1}V = \frac{(\delta^2 - \sigma^2)p^2}{2\mu\delta}$, where V is required to be a linear potential $V = \delta kx$, such as, the gravitational potential energy of a particle and the potential energy of an electric charge in an electric field of a uniform strength, which gives (15) for $k = 0$. As mentioned above, it is only classically acceptable for $\frac{\delta^2 - \sigma^2}{\delta} = 1$ with p and E being interpreted as momentum and kinetic energy, respectively.

In fact, as well known, not each effect of quantum mechanics can have the classical counterpart. So, the other cases of (15) could probably be physically meaningful results. In

the following, we will give some discussions of the relation between the kinetic energy and momentum of a particle. The purpose of doing this is to attempt to reveal some possible quantum effects that are banned by classical mechanics. Let now us give a simple analysis of Eq. (15). The two special cases of it, $E = 0$ and $E < 0$, are interesting, which correspond to $\delta^2 - \sigma^2 = 0$ and $\frac{\delta^2 - \sigma^2}{\delta} < 0$ with nonzero δ , respectively, where p is assumed to be real, nonzero. They exceed the range of classical mechanics and also cannot be obtained from the usual Schrödinger equation but are the consequences of the symmetrized Schrödinger equation indeed. The latter of these two cases may imply a phantom particle of mass μ , momentum p . The phantom energy is said to be a negative kinetic energy, it is allowed in condensate physics, cosmology and the current astronomical observation [11, 12, 13]. Since the phantom energy may exist in nature, a particle of the mass μ , nonzero momentum p and zero kinetic energy should also be acceptable. Our discussions of the relation between kinetic energy and momentum can be to some extent consistent with the conjecture [14, 15] that for a given classical action $S = \int dt \frac{m}{2} \dot{x}^2 - V(x)$ with m mass and x the position coordinate of a particle there is a quantum action $S = \int dt \frac{\tilde{m}}{2} \dot{x}^2 - \tilde{V}(x)$ in the path integral approach, where the kinetic energy and potential in the quantum action take the modified forms. In fact, this has also suggested that the relation between kinetic energy and momentum in quantum mechanics needn't to be equivalent completely to that in classical mechanics.

References

- [1] R. L. Liboff, *Introductory quantum mechanics*, (Holden-day, America, 1980).
- [2] E. Schrödinger, *Ann. Physik*, **79**, 489 (1926); **81**, 109 (1926).
- [3] M. Born, *Z. Physik*, **37**, 863 (1926); *Nature* **119**, 354 (1927).

- [4] F. Laloë, *Am. J. Phys.*, **69**, 655(2001).
- [5] M. Grigorescu, *Physical Framework of Quantization Problem*, arXiv:physics/0306079, (2003); *Rom. Rep. Phys.*, **45**, 645 (1993).
- [6] I. M. Yaglom, *Complex Numbers in Geometry*, Academic Press, London (1968).
- [7] Y. H. Wei, *Inter.J.Theor.Phys.*, **36**, 1617 (1997).
- [8] G. Kunsttater, J. W. Moffat and R. Malzan, *J. Math. Phys.*, **24**, 866(1983).
- [9] P. Bracken and J. Hayes, *Am. J. Phys.*, **71**, 726(2003).
- [10] Y. H. Wei, *Class. quantum Grav.*, **18**, 2163 (2001).
- [11] M. P. Dabrowski, T. Stachowiak and M. Szydłowski, arXiv:hep-ph/0307128.
- [12] R. R. Caldwell, M. Kamionkowski, and N. N. Weinberg *Phys. Rev. Lett.*, **91**, 071301 (2003); astro-ph/0302506.
- [13] R. R. Caldwell, *Phys. Lett. B* **545**, 23 (2002); astro-ph/9908168.
- [14] H. Jirari, H. Kröger, X. Q. Luo, K. J. M. Moriarty, and S. G. Rubin *Phys. Rev. Lett.*, **86**, 187 (2001).
- [15] H. Kröger, *Phys. Rev. A*, **65**, 052118 (2002).